

**Blake M. Bodendorfer, MD**  
**1 Edmundson Place Suite 500**  
**Council Bluffs, IA 51503**  
**712-323-5333**  
**16221 Evans Plaza**  
**Omaha, NE 68116**  
**402-991-9958**  
[bbodendorfer@millerortho.com](mailto:bbodendorfer@millerortho.com)  
[omahasportsdoc.com](http://omahasportsdoc.com)



### **Post-Operative Rehabilitation Guidelines for Distal Biceps Tendon Repair**

0-First Post-Op Visit: Sling and Splint all times

2-6 Weeks: Please make patient removable extension block splint at 45°  
Begin Active Assist ROM. Initially Limit Extension to 45°, Advance to full extension by 6 weeks.  
Splint all times other than exercises  
Wrist/Shoulder ROM exercises

6-9 Weeks: May D/C elbow Splint  
Continue Passive/Active Assist ROM Elbow  
Begin Biceps Isometrics  
Begin Cuff/Deltoid Isometrics  
Maintain Wrist/Shoulder ROM

9-12 Weeks: Begin Active Elbow Flexion against gravity  
Maintain ROM Elbow/Wrist/Shoulder  
Advance to resistive Strengthening Deltoid/Rotator Cuff

Week 12 – 6 months: Begin Gradual elbow flexion strengthening and advance as tolerated.